

Universality of Random Matrices and Local Relaxation Flow

László Erdős^{1*}, Benjamin Schlein² and Horng-Tzer Yau^{3†}

Institute of Mathematics, University of Munich,
Theresienstr. 39, D-80333 Munich, Germany¹

Department of Pure Mathematics and Mathematical Statistics
University of Cambridge
Wilberforce Rd, Cambridge CB3 0WB, UK²

Department of Mathematics, Harvard University
Cambridge MA 02138, USA³

Oct 28, 2009

Abstract

We consider $N \times N$ symmetric random matrices where the probability distribution for each matrix element is given by a measure ν with a subexponential decay. We prove that the eigenvalue spacing statistics in the bulk of the spectrum for these matrices and for GOE are the same in the limit $N \rightarrow \infty$. Our approach is based on the study of the Dyson Brownian motion via a related new dynamics, the local relaxation flow.

AMS Subject Classification: 15A52, 82B44

Running title: Universality for Wigner matrices

Key words: Wigner random matrix, Dyson Brownian Motion.

1 Introduction

A central question concerning random matrices is the universality conjecture which states that local statistics of eigenvalues are determined by the symmetries of the ensembles but are otherwise independent of the details of the distributions. There are two types of universalities: the edge universality and the bulk universality concerning the interior of the spectrum. The edge universality is commonly approached via the moment

*Partially supported by SFB-TR 12 Grant of the German Research Council

†Partially supported by NSF grants DMS-0757425, 0804279

method [20, 21] while the bulk universality was proven for very general classes of unitary invariant ensembles (see, e.g. [3, 5, 6, 18, 19] and references therein) based on detailed analysis of orthogonal polynomials. For non-unitary ensembles, recent results by Tao-Vu and the current authors [11, 22, 12] have established the bulk universality for *Hermitian* Wigner ensembles. The approaches of [11] and [22] are based on the following two common ingredients: (1) Johansson's [16] result (see also [2]) stating that bulk universality holds for ensembles of the form

$$\hat{H} + sV, \quad (1.1)$$

where $\hat{H}$ is a Wigner matrix, V is an independent standard GUE matrix and s is a positive constant. (2) For a random matrix H , find a random matrix $\hat{H}$ so that the eigenvalue statistics between H and $\hat{H} + sV$ are near each other.

Johansson's proof relied on the asymptotic analysis of an *explicit* formula by Brézin-Hikami [4, 16] for the correlation functions of the eigenvalues of $\hat{H} + sV$. Unfortunately, the similar formula for GOE is not very explicit and the corresponding result is not available. On the other hand, if we vary the parameter s in the matrix $\hat{H} + sV$ and s^2 is interpreted as time, then the evolution of the eigenvalues is the Dyson Brownian motion (DBM) [7]. If we replace the Brownian motions by the Ornstein-Uhlenbeck processes, the resulting dynamics on the eigenvalues, which we still call DBM, has the GUE eigenvalue distribution as the invariant measure. Thus the result of Johansson can be interpreted as stating that the local statistics of GUE is reached via DBM for time of order one. In fact, by analyzing the dynamics of DBM with ideas from the hydrodynamical limit, we have extended Johansson's result to $s^2 \gg N^{-3/4}$ [10]. The key observation of [10] is that the local statistics of eigenvalues depend exclusively on the approach to local equilibrium. This approach avoids the usage of explicit formulae for correlation functions, but the identification of local equilibria, unfortunately, still uses explicit representations of correlation functions by orthogonal polynomials (following e.g. [19]), and the extension to other ensembles is not a simple task.

Therefore, the universality for symmetric random matrices remained open and the only partial result is given by Tao-Vu (Theorem 23 in [22]) for Wigner matrices with the first four moments of the matrix elements matching those of GOE. In this paper, we introduce a dynamical approach based on a new flow, the local relaxation flow, which locally behaves like DBM, but has a faster decay to equilibrium. In this approach the analysis of orthogonal polynomials or explicit formulae are completely eliminated and the method applies to both Hermitian and symmetric ensembles. In fact, the heart of the proof is a convex analysis and it applies to β -ensembles for any $\beta \geq 1$. The model specific information required to complete this approach involves only rough estimates on the accuracy of the local density of states. We expect this method will apply to a very general class of models.

2 Statement of Main Results

To fix the notation, we will present the case of symmetric Wigner matrices; the modification to the Hermitian case is straightforward and will be omitted. Fix $N \in \mathbb{N}$ and we consider a symmetric matrix ensemble of $N \times N$ matrices $H = (h_{\ell k})$ with normalization given by

$$h_{\ell k} = N^{-1/2} x_{\ell k}, \quad (2.1)$$

where $x_{\ell k}$ for $\ell < k$ are independent, identically distributed random variables with the distribution ν that has zero expectation and variance 1. The diagonal elements $x_{\ell \ell}$ are also i.i.d. with distribution $\tilde{\nu}$ that has zero expectation and variance two.

The joint distributions of the eigenvalues $\mathbf{x} = (x_1, x_2, \dots, x_N)$ of the Gaussian Unitary Ensemble (GUE) and the Gaussian Orthogonal Ensemble (GOE) are given by the following measure

$$\mu = \mu_N(d\mathbf{x}) = \frac{e^{-\mathcal{H}(\mathbf{x})}}{Z_\beta} d\mathbf{x}, \quad \mathcal{H}(\mathbf{x}) = N \left[\beta \sum_{i=1}^N \frac{x_i^2}{4} - \frac{\beta}{N} \sum_{i < j} \log |x_j - x_i| \right] \quad (2.2)$$

where $\beta = 1$ for GOE and $\beta = 2$ for GUE. We consider μ defined on the set of *ordered* eigenvalues

$$\Sigma_N := \{\mathbf{x} \in \mathbb{R}^N : x_1 < x_2 < \dots < x_N\}.$$

For definiteness, consider the $\beta = 1$ GOE case. Suppose the matrix elements evolve according to the Ornstein-Uhlenbeck (OU) process on $\mathbb{R}$, i.e. the density of their distribution $\nu_t = u_t(x)dx$ satisfies

$$\partial_t u_t = \mathcal{L} u_t, \quad \mathcal{L} = \frac{1}{2} \frac{\partial^2}{\partial x^2} - \frac{x}{2} \frac{\partial}{\partial x} \quad (2.3)$$

with the reversible measure $e^{-x^2/2}dx$ (strictly speaking, a differently normalized OU process is used for the diagonal elements but we omit this detail here). The Ornstein-Uhlenbeck process (2.3) induces a stochastic process, the Dyson Brownian motion, on the eigenvalues with a generator given by

$$L = \sum_{i=1}^N \frac{1}{2N} \partial_i^2 + \sum_{i=1}^N \left(-\frac{\beta}{4} x_i + \frac{\beta}{2N} \sum_{j \neq i} \frac{1}{x_i - x_j} \right) \partial_i \quad (2.4)$$

acting on $L^2(\mu)$. Let

$$D(f) = - \int f L f d\mu = \sum_{j=1}^N \frac{1}{2N} \int (\partial_j f)^2 d\mu \quad (2.5)$$

be the corresponding Dirichlet form, where $\partial_j = \partial_{x_j}$. Clearly μ is a reversible measure for the dynamics generated by L . Denote the distribution of the eigenvalues at time t by $f_t(\mathbf{x})\mu(d\mathbf{x})$. Then f_t satisfies

$$\partial_t f_t = L f_t. \quad (2.6)$$

The corresponding stochastic differential equation for the eigenvalues $\mathbf{x}(t)$ is now given by (see, e.g. Section 12.1 of [14])

$$dx_i = \frac{dB_i}{\sqrt{N}} + \left[-\frac{\beta}{4} x_i + \frac{\beta}{2N} \sum_{j \neq i} \frac{1}{x_i - x_j} \right] dt, \quad 1 \leq i \leq N, \quad (2.7)$$

where $\{B_i : 1 \leq i \leq N\}$ is a collection of independent Brownian motions. The well-posedness of DBM has been proved in Section 4.3.1 of [14], see the Appendix for more details.

Note that the equations (2.6) and (2.7) are defined for any $\beta \geq 1$, independently of the original matrix models. Our main Theorem 2.1 will be stated for general $\beta \geq 1$.

Convention. We will use the letters C and c to denote general constants whose precise values are irrelevant and they may change from line to line.

2.1 Local Relaxation Flow

We now introduce the local relaxation measure, which has the local statistics of GOE (or GUE) but has a faster decaying dynamics. Recall that

$$\varrho_{sc}(x) := \frac{1}{2\pi} \sqrt{(4 - x^2)_+}$$

is the density of the semicircle law. Define

$$n_{sc}(E) := \int_{-\infty}^E \varrho_{sc}(x) dx, \quad (2.8)$$

and let

$$\gamma_j = n_{sc}^{-1}(j/N). \quad (2.9)$$

Fix a positive number η with $N^{-1/6} \ll \eta \ll 1$ and for the rest of this paper, $\varepsilon > 0$ is a small positive number which we will not specify. Let $\gamma_j^\pm := \gamma_j \pm \eta N^{-\varepsilon}$ and define the mean field potential of eigenvalues far away from the j -th one as

$$W_j(x) = -\frac{\beta}{N} \sum_{k: |k-j| \geq N\eta} \log(|x - \gamma_k| + \eta) \quad \text{if } x \in I_j := (\gamma_j^-, \gamma_j^+). \quad (2.10)$$

For $x \geq \gamma_j^+$, we extend W_j by

$$W_j(x) = W_j(\gamma_j^+) + W_j'(\gamma_j^+) (x - \gamma_j^+) + \frac{W_j''(\gamma_j^+)}{2} (x - \gamma_j^+)^2 \quad (2.11)$$

and similarly for $x \leq \gamma_j^-$. In other words, W_j is just the simplest convex extension of the function defined by (2.10) on I_j . This modification will avoid the singularities at $x = \gamma_k$. Notice that this is purely a technical device since we will show in (4.17) of Proposition 4.4 that the regime I_j^ε is negligible in the sense that

$$\sum_{j=1}^N \int f_t \mathbf{1}(x_j \notin I_j) d\mu \leq C e^{-cN^\varepsilon}. \quad (2.12)$$

The local relaxation measure $\omega_N = \omega$ is defined by

$$\omega = \frac{1}{\tilde{Z}} \exp \left[-N \sum_{j=1}^N \left\{ \beta \frac{x_j^2}{4} + W_j(x_j) \right\} + \beta \sum_{i < j} \log |x_i - x_j| - \frac{\beta}{2} \sum_i \sum_{j: |j-i| > N\eta} \log(|x_i - x_j| + \eta) \right] =: \frac{e^{-\tilde{H}}}{\tilde{Z}}$$

Let

$$\omega = \psi \mu, \quad \psi = \frac{Z_\beta}{\tilde{Z}} \exp \left[-\frac{\beta}{2} \sum_{i=1}^N \sum_{j: |j-i| > N\eta} \log(|x_i - x_j| + \eta) - N \sum_{i=1}^N W_i(x_i) \right].$$

Recall the definition of the relative entropy with respect to a measure λ

$$S_\lambda(f) = \int f \log f d\lambda, \quad S_\lambda(f|\psi) = \int f \log(f/\psi) d\lambda$$

and the Dirichlet form

$$D_\lambda(h) = \frac{1}{2N} \int (\nabla h)^2 d\lambda = \frac{1}{2N} \sum_j \int (\partial_j h)^2 d\lambda. \quad (2.13)$$

The local relaxation flow is defined to be the reversible dynamics w.r.t. ω characterized by the generator $\tilde{L}$ defined by

$$\int f \tilde{L} g d\omega = -\frac{1}{2N} \sum_j \int \partial_j f \partial_j g d\omega \quad (2.14)$$

Explicitly, $\tilde{L}$ is given by

$$L = \tilde{L} + \sum_j b_j \partial_j, \quad b_j = \frac{1}{N} \sum_{k: |k-j| > N\eta} \frac{\text{sgn}(x_j - x_k)}{|x_j - x_k| + \eta} + W'_j(x_j), \quad (2.15)$$

where

$$W'_j(x) = -\frac{1}{N} \sum_{k: |k-j| > N\eta} \frac{\text{sgn}(x - \gamma_k)}{|x - \gamma_k| + \eta} \quad (2.16)$$

for $x \in I_j$. Note that for any k with $|k-j| > N\eta$, we have $\gamma_k \notin 2I_j$, where $2I_j$ is the doubling of the interval I_j . Moreover, for $k = j \pm N\eta$ we have $|\gamma_k - \gamma_j| \leq C\eta^{2/3}$ with some constant C , so $|x - \gamma_k| \leq C\eta^{2/3}$ for $x \in I_j$. Thus we obtain that

$$\inf_{x \in I_j} W''_j(x) \geq c \inf_{|x| \leq 2+\eta} \int_{|x-\gamma| \geq C\eta^{2/3}} \frac{\varrho_{sc}(\gamma) d\gamma}{|x - \gamma|^2} \geq c\eta^{-1/3}.$$

Because W_j was defined by a convex extension outside of I_j , the same bound holds for any x :

$$\inf_j \inf_{x \in \mathbb{R}} W''_j(x) \geq c\eta^{-1/3}, \quad (2.17)$$

i.e., the mean field potential is uniformly convex.

We can now state the main result.

Theorem 2.1 (Universality of Dyson Brownian Motion for Short Time) *Suppose that $S_\mu(f_0|\psi) \leq CN^m$ for some m fixed. Let $\tau = \eta^{1/3}N^\varepsilon$ with some $\varepsilon > 0$ and assume that $\eta \geq N^{-1/33+\varepsilon}$. Assume that there is a positive number Λ such that*

$$\sup_{t \leq \tau} N \sum_j \int b_j^2 f_t d\mu \leq C\eta^{-2}\Lambda. \quad (2.18)$$

Fix $n \geq 1$, let $G : \mathbb{R}^n \rightarrow \mathbb{R}$ be a bounded smooth function with compact support and define

$$\mathcal{G}_{i,n}(\mathbf{x}) := G\left(N(x_i - x_{i+1}), N(x_{i+1} - x_{i+2}), \dots, N(x_{i+n-1} - x_{i+n})\right). \quad (2.19)$$

Set $\tau \geq R^2 N^\varepsilon$. Then for any $J \subset \{1, 2, \dots, N-n\}$ we have

$$\left| \int \frac{1}{N} \sum_{i \in J} \mathcal{G}_{i,n}(\mathbf{x}) f_\tau d\mu - \int \frac{1}{N} \sum_{i \in J} \mathcal{G}_{i,n}(\mathbf{x}) d\mu \right| \leq CN^{-1/2+\varepsilon/2} \eta^{-5/6} \Lambda^{1/2}. \quad (2.20)$$

We emphasize that Theorem 2.1 applies to *all* $\beta \geq 1$ *ensembles* and the only assumption concerning the distribution f_τ is in (2.18). The potential W is chosen to satisfy the two convexity properties: (2.17) and (3.7). Potentially, there are many other choices for W . For example, without changing the form of W given in (2.10), a more natural choice for γ_j would be

$$\gamma_j = \int x_j f_0 d\mu.$$

This would somewhat improve the estimate (2.22), but the analysis is more complicated and we will not pursue this choice in this paper.

Returning to the original Wigner ensembles with $\beta = 1$, we can estimate the critical constant Λ in the following lemma, to be proved in Section 4. We assume that the measure $d\nu$ satisfies the logarithmic Sobolev inequality (LSI), i.e. there is a constant θ such that for any nonnegative function u with $\int u d\nu = 1$ we have

$$\int u \log u d\nu \leq \theta \int |\nabla \sqrt{u}|^2 d\nu. \quad (2.21)$$

We remark that (2.21) implies that ν has a Gaussian decay.

Lemma 2.2 *Suppose the assumption (2.21) on the distribution ν of the matrix elements hold. Then the constant Λ in (2.18) can be estimated as*

$$\Lambda \leq C_\sigma \eta^{-2} N^{8/9+\sigma} \quad (2.22)$$

for any $\sigma > 0$.

Now we state the universality of the local eigenvalue statistics for Wigner matrices. To formulate the result in terms of correlation functions, it is convenient to remove the ordering $x_1 < x_2 < \dots < x_N$ among the eigenvalues and work with symmetric densities. Let $p_N(x_1, x_2, \dots, x_N)$ denote the (symmetric) probability density of eigenvalues and for any $k = 1, 2, \dots, N$ let

$$p_N^{(k)}(x_1, x_2, \dots, x_k) := \int_{\mathbb{R}^{N-k}} p_N(x_1, x_2, \dots, x_N) dx_{k+1} \dots dx_N \quad (2.23)$$

be the k -point correlation function. In the case of GOE, we denote the density functions by $p_{N,GOE}^{(k)}$ and it is well known that for $|u| < 2$ the limit

$$\frac{1}{[\varrho_{sc}(u)]^k} p_{N,GOE}^{(k)}\left(u + \frac{\alpha_1}{N\varrho_{sc}(u)}, \dots, u + \frac{\alpha_k}{N\varrho_{sc}(u)}\right) \quad (2.24)$$

exists and can be explicitly computed (see, e.g. Section 7 of [18]).

Theorem 2.3 *Suppose the distribution ν for the matrix elements satisfies the assumptions (2.21). Assume that the density $\nu(x) = e^{-U(x)}$ satisfies*

$$\sum_{j=1}^m |U^{(j)}(x)| \leq C(1+x^2)^\ell \quad (2.25)$$

for some $\ell \in \mathbb{N}$ and $m \in \mathbb{N}$ sufficiently large. Let $k \geq 1$ and $O : \mathbb{R}^k \rightarrow \mathbb{R}$ be a continuous, compactly supported function. Then for any $|u| < 2$, we have

$$\begin{aligned} \lim_{\delta \rightarrow 0} \lim_{N \rightarrow \infty} \frac{1}{2\delta} \int_{u-\delta}^{u+\delta} \int_{\mathbb{R}^k} O(\alpha_1, \dots, \alpha_k) \\ \times \frac{1}{[\varrho_{sc}(v)]^k} \left(p_N^{(k)} - p_{N,GOE}^{(k)} \right) \left(v + \frac{\alpha_1}{N\varrho_{sc}(v)}, \dots, v + \frac{\alpha_k}{N\varrho_{sc}(v)} \right) d\alpha_1 \dots d\alpha_k dv = 0. \end{aligned} \quad (2.26)$$

In the following Corollary we remove all assumption from Theorem 2.3 except for the decay condition and a technical condition that ν is supported in at least three points. This latter technical condition was removed in our later paper [13], where we generalized our approach to a broader class of random matrix ensembles.

Corollary 2.4 *Suppose the distribution ν of the matrix elements has mean zero, variance one and a tail with subexponential decay, i.e. it satisfies that*

$$\int e^{|x|^c} d\nu(x) < \infty \quad (2.27)$$

for some constants $c > 0$. Assume that ν is supported in at least three points. Then the conclusion of Theorem 2.3 holds.

Proof. Let m_j denote the moments of ν

$$m_j := \int x^j d\nu,$$

where $m_1 = 0$ and $m_2 = 1$. It is easy to check that $m_4 \geq m_3^2 + 1$ with the equality holds only for Bernoulli type distribution supported in two points. Let $\widehat{\nu}$ be the probability measure on $\mathbb{R}$ with density

$$e^{-U(x)} = (\text{const.})e^{-x^2/2+F(x)}$$

where F is a smooth function with compact support. For the fixed numbers m_3, m_4 satisfying $m_4 > m_3^2 + 1$, there exists an F such that the first four moments of $\widehat{\nu}$ match to those of ν , i.e.

$$\int_{\mathbb{R}} x d\widehat{\nu} = 0, \quad \int_{\mathbb{R}} x^2 d\widehat{\nu} = 1, \quad \int_{\mathbb{R}} x^j d\widehat{\nu} = m_j, \quad j = 3, 4.$$

Here $\|F\|_{\infty}$ and the support of F depend only on m_3, m_4 . Clearly, $\widehat{\nu}$ satisfies the assumption of Theorem 2.3 and thus (2.26) holds for the measure $\widehat{\nu}$. Recall that Theorem 15 in [22] states that the local eigenvalue statistics for matrices whose matrix element distributions match up to the first four moments are the same in the limit $N \rightarrow \infty$ (strictly speaking, this theorem was stated only for hermitian matrices, but the parallel version for the symmetric matrix holds as well, see the remark at the end of Section 1.6 [22]). This proves the Corollary. $\square$

Theorem 2.3 is a simple corollary of Theorem 2.1, Lemma 2.2 and the method of the reverse heat flow [11]. Lemma 2.2 is a technical result concerning some rough estimates of eigenvalues. Though we stated the

universality in terms of correlation functions, it also holds for the eigenvalue gap distribution and we omit the obvious statement (the analogous statement for the Hermitian case was formulated in Theorem 1.2 of [11]).

We now sketch the key new ideas of this article.

- I. The key concept is the introduction of the local relaxation flow (2.14) which has the following two properties: (1) The invariant measure for this flow, the local relaxation measure ω will be shown to have the same local eigenvalue statistics as the GOE. (2) The relaxation time of the local relaxation flow is much shorter than that of the DBM, which is of order one.
- II. Suppose we have a density q w.r.t. ω . Then, by differentiating the Dirichlet form w.r.t. ω under the local relaxation flow, we will prove that the difference between the local statistics of $q\omega$ and ω can be estimated in terms of the Dirichlet form of q w.r.t. ω . Hence if the Dirichlet form is small, the local statistics of $q\omega$ is independent of q .
- III. It remains to show that the Dirichlet form of $q = f_t\mu$ w.r.t. ω is small for t sufficiently large (but still much less than order one). To do that, we study the evolution of the entropy of $f_t\mu$ relative to ω . This provides estimates on the entropy and Dirichlet form which serve as inputs for the Step II to conclude the universality.

3 Local Relaxation Flows and Entropy Dissipation Estimates

The first ingredient to prove Theorem 2.1 is the analysis of the local relaxation flow which satisfies the logarithmic Sobolev inequality and the following entropy dissipation estimate.

Theorem 3.1 (Dirichlet Form Dissipation Estimate) *Suppose (2.17) holds. Consider the equation*

$$\partial_t q_t = \tilde{L}q_t \tag{3.1}$$

with reversible measure ω . Then we have

$$\partial_t D_\omega(\sqrt{q_t}) \leq -C\eta^{-1/3} D_\omega(\sqrt{q_t}) - \frac{1}{2N^2} \int \sum_{|i-j| \leq N\eta} \frac{1}{(x_i - x_j)^2} (\partial_t \sqrt{q_t} - \partial_j \sqrt{q_t})^2 d\omega, \tag{3.2}$$

$$\frac{1}{2N^2} \int_0^\infty ds \int \sum_{|i-j| \leq N\eta} \frac{1}{(x_i - x_j)^2} (\partial_i \sqrt{q_s} - \partial_j \sqrt{q_s})^2 d\omega \leq D_\omega(\sqrt{q_0}) \tag{3.3}$$

and the logarithmic Sobolev inequality

$$S_\omega(q) \leq C\eta^{1/3} D_\omega(\sqrt{q}) \tag{3.4}$$

with a universal constant C . Thus the time to equilibrium is of order $\eta^{1/3}$ and we have

$$S_\omega(q_t) \leq e^{-Ct\eta^{-1/3}} S_\omega(q_0). \tag{3.5}$$

The proof given below follows the argument in [1] and it was outlined in this context in [10]. The new observation is the additional second term on the r.h.s of (3.2), corresponding to “local Dirichlet form dissipation”. The estimate (3.3) on this additional term will play a key role in this paper.

Proof. Recall that from the derivation of the LSI (see, e.g. Section 5.1 of [10]), with $h = \sqrt{q}$ we have

$$\partial_t h = \tilde{L}h + \frac{1}{2N}h^{-1}(\nabla h)^2$$

and

$$\partial_t \frac{1}{2N} \int (\nabla h)^2 e^{-\tilde{\mathcal{H}}} d\mathbf{x} \leq -\frac{1}{2N^2} \int \nabla h (\nabla^2 \tilde{\mathcal{H}}) \nabla h e^{-\tilde{\mathcal{H}}} d\mathbf{x}. \quad (3.6)$$

In our case, (2.17) and the fact that

$$\frac{d^2}{dx^2} \left(\log(|x| + \eta) - \log |x| \right) \geq 0, \quad x > 0,$$

imply that the Hessian of $\tilde{\mathcal{H}}$ is bounded from below as

$$\frac{1}{2N^2} \nabla h (\nabla^2 \tilde{\mathcal{H}}) \nabla h \geq C\eta^{-1/3} \frac{1}{N} \sum_j (\partial_j h)^2 + \frac{1}{2N^2} \sum_{|i-j| \leq N\eta} \frac{1}{(x_i - x_j)^2} (\partial_i h - \partial_j h)^2 \quad (3.7)$$

with some positive constant C . This proves (3.2) and (3.3). Inserting the inequality

$$\partial_t D_\omega(\sqrt{q_t}) \leq -C\eta^{-1/3} D_\omega(\sqrt{q_t})$$

from (3.2) into the equation

$$\partial_t S_\omega(q_t) = -4D_\omega(\sqrt{q_t}), \quad (3.8)$$

and integrating the resulting equation, we prove (3.4). Inserting (3.4) into (3.8) we have

$$\partial_t S_\omega(q_t) \leq -C\eta^{-1/3} S_\omega(q_t)$$

and we obtain (3.5). □

Remark 3.2 *The proof of (3.6) requires an integration by parts and the boundary term at $x_i = x_j$ (explained in Section 5.1. of [10]) should vanish. In the Appendix we will justify this technical step.*

Lemma 3.3 *Suppose that the density q_0 satisfies $S_\omega(q_0) \leq CN^m$ with some $m > 0$ fixed. For a fixed $n \geq 1$ let $G : \mathbb{R}^n \rightarrow \mathbb{R}$ be a bounded smooth function with compact support and recall the definition of $\mathcal{G}_{i,n}$ from (2.19). Let $J \subset \{1, 2, \dots, N - n\}$ and set $\tau = \eta^{1/3} N^\varepsilon$. Then we have*

$$\left| \int \frac{1}{N} \sum_{i \in J} \mathcal{G}_{i,n}(\mathbf{x}) d\omega - \int \frac{1}{N} \sum_{i \in J} \mathcal{G}_{i,n}(\mathbf{x}) q_0 d\omega \right| \leq C \sqrt{\frac{D_\omega(\sqrt{q_0})^\tau}{N}} + Ce^{-cN^\varepsilon} \quad (3.9)$$

with some constant C depending on G .

Proof. Without loss of generality, we consider only the case $J = \{1, \dots, N\}$. Let q_t satisfy

$$\partial_t q_t = \tilde{L} q_t$$

with an initial condition q_0 . We first compare q_τ with $q_\infty = 1$. Using the entropy inequality

$$\int |q - 1| d\omega \leq \sqrt{S_\omega(q)}$$

and the exponential decay of the entropy (3.5), we have

$$\left| \int \frac{1}{N} \sum_i \mathcal{G}_{i,n}(\mathbf{x}) q_\tau d\omega - \int \frac{1}{N} \sum_i \mathcal{G}_{i,n}(\mathbf{x}) d\omega \right| \leq C(N^m e^{-\tau \eta^{-1/3}})^{1/2} \leq C e^{-cN^\varepsilon}.$$

To compare q_0 with q_τ , by differentiation, we have

$$\begin{aligned} & \int \frac{1}{N} \sum_i \mathcal{G}_{i,n}(\mathbf{x}) q_\tau d\omega - \int \frac{1}{N} \sum_i \mathcal{G}_{i,n}(\mathbf{x}) q_0 d\omega \\ &= \int_0^\tau ds \int \frac{1}{N} \sum_i \sum_{k=1}^n \partial_k G(N(x_i - x_{i+1}), \dots, N(x_{i+n-1} - x_{i+n})) [\partial_{i+k-1} q_s - \partial_{i+k} q_s] d\omega. \end{aligned}$$

From the Schwarz inequality and $\partial q = 2\sqrt{q}\partial\sqrt{q}$ the last term is bounded by

$$\begin{aligned} & 2 \sum_{k=1}^n \left[\int_0^\tau ds \int \sum_i \left[\partial_k G(N(x_i - x_{i+1}), \dots, N(x_{i+n-1} - x_{i+n})) \right]^2 (x_{i+k-1} - x_{i+k})^2 q_s d\omega \right]^{1/2} \\ & \times \left[\int_0^\tau ds \int \frac{1}{N^2} \sum_i \frac{1}{(x_{i+k-1} - x_{i+k})^2} [\partial_{i+k-1} \sqrt{q_s} - \partial_{i+k} \sqrt{q_s}]^2 d\omega \right]^{1/2} \\ & \leq C_n \sqrt{\frac{D_\omega(\sqrt{q_0})\tau}{N}}, \end{aligned} \tag{3.10}$$

where we have used (3.3) and that

$$\left[\partial_k G(N(x_i - x_{i+1}), \dots, N(x_{i+k-1} - x_{i+k}), \dots, N(x_{i+n-1} - x_{i+n})) \right]^2 (x_{i+k-1} - x_{i+k})^2 \leq C N^{-2},$$

since G is smooth and compactly supported. This proves the Lemma. $\square$

Notice if we use only the entropy dissipation and Dirichlet form, the main term on the right hand side of (3.9) will become $\sqrt{S}\tau$. Hence by exploiting the Dirichlet form dissipation coming from the second term on the r.h.s. of (3.2), we gain the crucial factor $N^{-1/2}$ in the estimate.

The second ingredient to prove Theorem 2.1 is the following entropy and Dirichlet form estimates. The first estimate (3.11) should be compared with the bound $S_\mu(f_t) \leq C N^{5/4}$ for $t \geq N^{-1}$ given in Lemma 5.1 of [10].

Theorem 3.4 (Entropy and Dirichlet Form Estimates) *Suppose the assumptions of Theorem 2.1 hold. Let $\tau = \eta^{1/3} N^\varepsilon$ and let $g_t = f_t/\psi$ so that $S_\mu(f_t|\psi) = S_\omega(g_t)$. Then the entropy and the Dirichlet form satisfy the estimates:*

$$S_\omega(g_{\tau/2}) \leq C\eta^{-5/3}\Lambda, \quad (3.11)$$

$$D_\omega(\sqrt{g_\tau}) \leq 2\tau^{-1}S_\omega(g_{\tau/2}) + C\eta^{-2} \leq C\eta^{-2}\Lambda. \quad (3.12)$$

Proof. First we need the following relative entropy identity from [24].

Lemma 3.5 *Let f_t be a probability density satisfying $\partial_t f_t = Lf_t$. Then for any probability density ψ_t we have*

$$\partial_t S_\mu(f_t|\psi_t) = -\frac{2}{N} \sum_j \int (\partial_j \sqrt{g_t})^2 \psi_t d\mu + \int g_t (L - \partial_t) \psi_t d\mu,$$

where $g_t = f_t/\psi_t$.

In our setting, ψ is indep of t and L satisfies (2.15). Hence we have

$$\partial_t S_\omega(g_t) = \partial_t S_\mu(f_t|\psi) = -\frac{2}{N} \sum_j \int (\partial_j \sqrt{g_t})^2 d\omega + \int \tilde{L} g_t d\omega + \sum_j \int b_j \partial_j g_t d\omega.$$

Since the middle term on the right hand side vanishes, we have from the Schwarz inequality

$$\partial_t S_\omega(g_t) \leq -D_\omega(\sqrt{g_t}) + CN \sum_j \int b_j^2 g_t d\omega. \quad (3.13)$$

Together with the LSI (3.4) and (2.18), we have

$$\partial_t S_\omega(g_t) \leq -C\eta^{-1/3} S_\omega(g_t) + C\eta^{-2}\Lambda. \quad (3.14)$$

for $t \leq \tau$. Since $S_\omega(g_0) = S_\mu(f_0|\psi) \leq N^m$ and $\tau/2 \gg \eta^{1/3}$, the last inequality proves (3.11).

Integrating (3.13) from $t = \tau/2$ to $t = \tau$ and using the monotonicity of the Dirichlet form in time, we have proved (3.12) with the choice of τ . □

Proof of Theorem 2.1. Fix $\tau = \eta^{1/3} N^\varepsilon$ and let $q_0 = g_\tau = f_\tau/\psi$ with f satisfying the assumption of Theorem 3.4. Using (3.12), we have

$$\sqrt{\frac{D_\omega(\sqrt{q_0})\tau}{N}} \leq CN^{-1/2+\varepsilon/2} \eta^{-5/6} \Lambda^{1/2},$$

and from (3.9) we also have

$$\left| \int \frac{1}{N} \sum_i \mathcal{G}_{i,n}(\mathbf{x}) f_\tau d\mu - \int \frac{1}{N} \sum_i \mathcal{G}_{i,n}(\mathbf{x}) d\omega \right| \leq CN^{-1/2+\varepsilon/2} \eta^{-5/6} \Lambda^{1/2} + Ce^{-cN^\varepsilon}. \quad (3.15)$$

Clearly, equation (3.15) also holds for the special choice $f_0 = 1$ (for which $f_\tau = 1$), i.e. local statistics of μ and ω can be compared. Hence we can replace the measure ω in (3.15) by μ and this proves Theorem 2.1. □

Denote by

$$\alpha_j = \alpha_j(t) = \mathbb{E}_t x_j \quad (3.16)$$

the expected location of x_j w.r.t f_t . We now prove Theorem 2.3.

Proof of Theorem 2.3. With the choice $\Lambda = C_\varepsilon \eta^{-2} N^{8/9+\varepsilon}$ in (2.18) and recalling that $\eta \geq N^{-1/33+\varepsilon}$, we have

$$N^{-1/2+\varepsilon/2} \eta^{-5/6} \Lambda^{1/2} \ll 1.$$

Thus we have proved that

$$\lim_{N \rightarrow \infty} \left| \int \frac{1}{N} \sum_{i \in J} \mathcal{G}_{i,n}(\mathbf{x}) f_\tau d\mu - \int \frac{1}{N} \sum_{i \in J} \mathcal{G}_{i,n}(\mathbf{x}) d\mu \right| = 0. \quad (3.17)$$

We now prove (2.26) for $k = 2$; the other cases are similar. Since we integrate over v , it suffices to consider only observables of the form $O(\alpha_1 - \alpha_2)$ with O being bounded and compactly supported. Denote $\varrho_{sc}(u) = a$ and we have

$$\int_{u-\delta}^{u+\delta} \int_{\mathbb{R}} O(\alpha) p_N^{(2)} \left(v, v + \frac{\alpha}{N \varrho_{sc}(v)} \right) d\alpha = \frac{a}{N} \sum_{i: |x_i - u| \leq \delta} \sum_{j: |x_j - u| \leq \delta} \int f(\mathbf{x}) O[Na(x_i - x_j)] d\mu + o(1) \quad (3.18)$$

where $o(1) \rightarrow 0$ as $N \rightarrow \infty$, $\delta \rightarrow 0$ and $f d\mu$ is the probability distribution of the eigenvalues of the random matrix.

Proposition 4.4 from Section 4 asserts that $\max_m |\alpha_m(t) - \gamma_m| \rightarrow 0$ as $N \rightarrow \infty$ both for $t = 0$ and $t = \tau$. Thus we can replace the conditions $|x_k - u| \leq \delta$ for $k = i, j$ in the summations with the condition $k \in J = [N n_{sc}(u - \delta), N n_{sc}(u + \delta)]$ at the expense of negligible errors. By using the method of Proposition 2.1 of [11] and that the derivatives on U are controlled by (2.25) for m large, we can find an initial data $\hat{f}$ such that the difference between the expectation on the right hand side of (3.18) w.r.t. f and $\hat{f}_\tau = e^{\tau L} \hat{f}$ is negligible. The distribution $\hat{f}$ can be found by approximating the reverse heat flow $e^{-\tau L} f$ by a sufficiently high order Taylor polynomial of $e^{-\tau L}$ as outlined in Remark 1.1 of [11].

Thus we only have to compare

$$\lim_{N \rightarrow \infty} \frac{a}{N} \sum_{i,j \in J} \int \hat{f}_\tau(\mathbf{x}) O[Na(x_i - x_j)] d\mu$$

with the GOE case. We can rewrite the last term as

$$\lim_{M \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{a}{N} \sum_{i \in J} \sum_{n \leq M} \int \hat{f}_\tau(\mathbf{x}) O[Na(x_i - x_{i+n})] d\mu. \quad (3.19)$$

The observable in (3.19) is of the form $\mathcal{G}_{i,n}(\mathbf{x})$ for an appropriate G . From Theorem 2.1, the limit (3.19) is unchanged if we replace $\hat{f}_\tau d\mu$ by $d\mu$. Here the assumptions of Theorem 2.1 w.r.t. $\hat{f}$ can be easily checked from the explicit construction in [11]. Note that the entropy assumption in Theorem 2.1 follows from the LSI with $m = 2$. This proves Theorem 2.3. $\square$

4 Proof of Lemma 2.2

Let $\mathbb{P}_t$ and $\mathbb{E}_t$ denote the probability and expectation with respect to the measure $f_t\mu$. We first collect the assumptions we need in this section.

1. Logarithmic Sobolev inequality (LSI) for the distribution ν_t of the matrix elements. Since LSI holds for ν (2.21), it also holds for ν_t as well; see Lemma B.1 in Appendix B for a proof.
2. Upper bound on the number of eigenvalues $\mathcal{N}_I$ in an interval of length $|I| \geq (\log N)^2/N$:

$$\mathbb{P}_t(\mathcal{N}_I \geq KN|I|) \leq Ce^{-c\sqrt{KN|I|}} \quad (4.1)$$

for any $K \geq K_0$ with a sufficiently large fixed K_0 . This bound was proved in Theorem 4.6 of [9] under the condition of Gaussian decay that follows from LSI (2.21). The conditions $y > (\log N)/N$ and $|I| \geq (\log N)/N$ were unfortunately erroneously stated in Theorem 4.6 of [9]; the proof given there requires that $y > (\log N)^2/N$ and $|I| \geq (\log N)^2/N$. Under a stronger condition on ν (see remark after Lemma 4.7 of [9]), the statement of Theorem 4.6 of [9] holds for $y > (\log N)/N$ and $|I| \geq (\log N)/N$.

3. (Theorem E1 of [10]) Let $|E| < 2$. Denote by x_α the largest eigenvalue below E and assume that $\alpha \leq N-1$. Then there exist positive constants C and c such that for any M that satisfy $c(\log N)^4/(2-|E|) \leq M \leq CN(2-|E|)$, we have

$$\mathbb{P}_t\left(x_{\alpha+1} - E \geq \frac{M}{N}, \alpha \leq N-1\right) \leq Ce^{-c[2-|E|]^{3/2}\sqrt{M}}. \quad (4.2)$$

4. (Equation (4.15) of [10]) Let

$$n(E) = \frac{1}{N}\mathbb{E}_t\#\{x_j \leq E\}. \quad (4.3)$$

Then there exists a constant $C > 0$ such that

$$\int_{-\infty}^{\infty} |n(E) - n_{sc}(E)|dE \leq \frac{C}{N^{6/7}}. \quad (4.4)$$

The real assumption behind these statements is LSI that was used in the proof of (4.2) and (4.4). Since the parameter t will not play any role in this section, we will drop the index from the notation, i.e. we use $\mathbb{E} = \mathbb{E}_t$, $\mathbb{P} = \mathbb{P}_t$ etc.

Proposition 4.4 below estimates the distance of the eigenvalues from their location given by the semicircle law. This will justify that the convex extension of the potential W_j affects only regimes of very small probability. Recall the definition (3.16) of $\alpha_j = \alpha_j(t)$ for the expected location of the j -th eigenvalue w.r.t $f_t\mu$. We start with an estimate on the expected location of the extreme eigenvalues:

Lemma 4.1 *Suppose that the probability measure ν of the matrix entries satisfies*

$$\int_{\mathbb{R}} e^{c|x|}d\nu(x) < \infty \quad (4.5)$$

for some $c > 0$ (this condition is satisfied, in particular, under LSI (2.21)). Then for any $\delta > 0$ we have

$$-2 - CN^{-1/4+\delta} \leq \alpha_1 < \alpha_N \leq 2 + CN^{-1/4+\delta} \quad (4.6)$$

with some C depending on δ .

Proof. For any M , define the probability measure ζ_M on $\mathbb{R}$ with density

$$Z_M^{-1} e^{-U(x)} 1(|x| \leq M).$$

Setting $M = N^{-\delta}$ and using (4.5), the total variational norm between ν and ζ_M is bounded

$$\|\zeta_M - \nu\|_\infty \leq C e^{-cN^\delta}.$$

Denote by $\zeta_M^N = \zeta_M^{\otimes N^2}$ (ν^N resp.) the probability law of the random matrices whose matrix elements are distributed according to ζ_M (ν resp.). Then the total variational norm between ζ_M^N and ν^N is bounded by

$$\|\zeta_M^N - \nu^N\|_\infty \leq C N^2 e^{-cN^\delta}.$$

Since up to exponential small probability, the largest eigenvalue x_N is bounded, the expectations of x_N w.r.t. these two measures satisfy

$$\left| \mathbb{E}_{\zeta_M^N} x_N - \mathbb{E}_{\nu^N} x_N \right| \leq C N^2 e^{-cN^\delta}.$$

From Theorem 1.4 of [23], we have

$$\mathbb{E}_{\zeta_M^N} x_N \leq 2 + C M^{1/2} N^{-1/4} \log N \leq 2 + N^{-1/4+\delta}$$

for any $\delta > 0$. Thus we have proved that

$$\alpha_N = \mathbb{E}_{\nu^N} x_N \leq 2 + C N^{-1/4+\delta}.$$

Similar estimate holds for α_1 . □

Next we estimate the fluctuations of x_j :

Proposition 4.2 *For any $\varepsilon > 0$ we have*

$$\mathbb{P}\left(\max_j |x_j - \alpha_j| \geq N^{-1/2+\varepsilon}\right) \leq C_\varepsilon e^{-cN^\varepsilon}. \quad (4.7)$$

Proof. Using (2.21) and the Bobkov-Götze concentration inequality, we have for any $T > 0$

$$\mathbb{P}\left(\max_j |x_j - \mathbb{E} x_j| \geq \gamma\right) \leq 2N \max_j \mathbb{P}\left(x_j - \mathbb{E} x_j \geq \gamma\right) \leq 2N e^{-\gamma T} \max_j \mathbb{E} e^{\theta T^2 |\nabla x_j|^2} \leq C N e^{-c\gamma^2 N}$$

after optimizing for T and using that $|\nabla x_j| \leq C N^{-1/2}$ (see the analogous calculation in the proof of Theorem 3.1 of [8]). This proves (4.7). □

The following proposition is a refinement of Proposition 4.2:

Proposition 4.3 *We choose M, K, κ positive numbers, depending on N , such that for some small $\delta > 0$*

$$N^{-1/6+2\delta} \leq \kappa \leq N^{-\delta}, \quad M \kappa^6 \geq N^{2\delta}, \quad K \leq c N \kappa^{3/2} \quad \text{and} \quad c(\log N)^4 / \kappa \leq M \leq C N \kappa \quad (4.8)$$

with some sufficiently small $c > 0$ and large $C > 0$ constants. Let

$$\Phi := 2N^{-1/2+2\delta} K^{-1/2} + \frac{2KM}{N}. \quad (4.9)$$

Then for any $CN\kappa^{3/2} \leq i \leq CN(1 - \kappa^{3/2})$ we have

$$\mathbb{P}\left(|x_i - \alpha_i| \geq \Phi\right) \leq Ce^{-cN^\delta}. \quad (4.10)$$

We also have

$$\mathbb{P}\left(\frac{1}{N} \sum_{i=1}^N |x_i - \alpha_i| \geq C\kappa^{3/2}N^{-1/2+\delta} + \Phi\right) \leq Ce^{-cN^\delta}. \quad (4.11)$$

The constants C may depend on δ but are independent of N .

Proof. Consider an index i with $CN\kappa^{3/2} \leq i \leq N(1 - C\kappa^{3/2})$, then $|2 - |\gamma_i|| \geq C\kappa$. We first show that $|2 - |x_i|| \geq C\kappa^2$ with a very high probability. Suppose, in the contrary, that $x_i < -2 + C\kappa^2$ for some $i \geq CN\kappa^{3/2}$ (the case $x_i \geq 2 - C\kappa^2$ is treated analogously). From (4.6) and (4.7) it follows that $x_1 \geq -2 - CN^{-1/4+\delta}$ with a very high probability. But then the interval $[-2 - CN^{-1/4+\delta}, -2 + C\kappa^2]$ of length $CN^{-1/4+\delta} + C\kappa^2 \ll \kappa^{3/2}$ would contain $CN\kappa^{3/2}$ eigenvalues, an event with an extremely low probability by (4.1).

Knowing that $|2 - |x_i|| \geq \kappa^2$ with a very high probability, we can use (4.2) to conclude that for any index i with $CN\kappa^{3/2} \leq i \leq N(1 - C\kappa^{3/2})$ we have

$$\mathbb{P}\left(x_{i+1} - x_i \geq \frac{M}{N}\right) \leq e^{-c\sqrt{M\kappa^6}} + Ce^{-cN^\delta} \leq Ce^{-cN^\delta} \quad (4.12)$$

by (4.8). Then

$$\mathbb{P}\left(\left|x_i - \frac{1}{2K+1} \sum_{|j-i| \leq K} x_j\right| \geq \frac{KM}{N}\right) \leq Ce^{-cN^\delta}. \quad (4.13)$$

Similarly to the calculation in Theorem 3.1 of [8], by using the logarithmic Sobolev inequality (2.21), we have

$$\mathbb{P}\left(\left|\frac{1}{2K+1} \sum_{j: |j-i| \leq K} x_j - \frac{1}{2K+1} \sum_{j: |j-i| \leq K} \mathbb{E}x_j\right| \geq N^{-1/2+2\delta}K^{-1/2}\right) \leq Ce^{-cN^\delta}. \quad (4.14)$$

This bound holds for any index i with the remark that if $i < K$ or $i > N - K$, then the averaging over the indices j is done asymmetrically.

Combining this estimate with (4.13) we have, apart from a set of very small probability, that

$$\left|x_i - \frac{1}{2K+1} \sum_{j: |j-i| \leq K} \alpha_j\right| \leq \Phi \quad (4.15)$$

for any $CN\kappa^{3/2} \leq i \leq N(1 - C\kappa^{3/2})$. Taking expectation, and using that $\mathbb{P}(\max_i |x_i| \geq \tilde{K}) \leq e^{-N\tilde{K}}$ for any sufficiently large $\tilde{K}$, we also obtain for these i indices that

$$\left|\alpha_i - \frac{1}{2K+1} \sum_{j: |j-i| \leq K} \alpha_j\right| \leq \Phi. \quad (4.16)$$

Subtracting the last two inequalities yields (4.10). Finally, combining this bound with the estimate (4.7) for the extreme indices, we obtain (4.11). $\square$

Proposition 4.4 *For any small $\delta > 0$ and for any $j = 1, 2, \dots, N$ we have*

$$\mathbb{P}\left(|x_j - \gamma_j| \geq CN^{-1/5+\delta}\right) \leq Ce^{-cN^\delta} \quad (4.17)$$

and we also have

$$\max_m |\alpha_m - \gamma_m| \leq CN^{-1/5+\delta}. \quad (4.18)$$

The constants C may depend on δ but are independent of N .

We remark that in the bulk $\alpha_m - \gamma_m$ is expected to be bounded by $O(N^{-1+\epsilon})$ (in the hermitian case it was proven in [15], see also [22]); near the edges one expects $\alpha_m - \gamma_m \sim O(N^{-2/3})$. Our estimate is not optimal, but it gives a short proof that is sufficient for our purpose.

Proof of Proposition 4.4. We define

$$\tilde{n}(E) := \frac{1}{N} \#\{\alpha_j \leq E\}$$

to be the counting function of the expected locations of the eigenvalues. We compare $\tilde{n}(E)$ with $n(E)$ defined in (4.3). Using the fluctuation bound (4.7), we have

$$n(E - N^{-1/2+\epsilon}) - Ce^{-cN^\epsilon} \leq \tilde{n}(E) \leq n(E + N^{-1/2+\epsilon}) + Ce^{-cN^\epsilon}$$

for any $E \in \mathbb{R}$. In fact, the upper bound on the density (4.1) guarantees that $\tilde{n}(E)$ and $n(E)$ are Lipschitz continuous on any scale much bigger than $(\log N)^2/N$, i.e.

$$|\tilde{n}(E) - n(E)| \leq CN^{-1/2+\epsilon} + Ce^{-cN^\epsilon} \leq CN^{-1/2+\epsilon} \quad (4.19)$$

for any E .

We write

$$\sum_{j=1}^N |\alpha_j - \gamma_j| = \sum_{j:\alpha_j \geq \gamma_j} (\alpha_j - \gamma_j) + \sum_{j:\alpha_j \leq \gamma_j} (\gamma_j - \alpha_j).$$

For the first term

$$\begin{aligned} \sum_{j:\alpha_j \geq \gamma_j} (\alpha_j - \gamma_j) &= \int dE \sum_{j:\alpha_j \geq \gamma_j} \mathbf{1}(\gamma_j \leq E \leq \alpha_j) = \int dE \sum_{j:\alpha_j \geq \gamma_j} \mathbf{1}\left(\tilde{n}(E) \geq \frac{j}{N} > n_{sc}(E)\right) \\ &= N \int dE \mathbf{1}(n_{sc}(E) \leq \tilde{n}(E)) (n_{sc}(E) - \tilde{n}(E)) \end{aligned} \quad (4.20)$$

and the second term is analogous. We thus have

$$\frac{1}{N} \sum_{j=1}^N |\alpha_j - \gamma_j| = \int |\tilde{n}(E) - n_{sc}(E)| dE \quad (4.21)$$

By Lemma 4.1

$$\int_{|E| \geq 3} |\tilde{n}(E) - n_{sc}(E)| dE = 0.$$

For the energy range $|E| \leq 3$, we use (4.19):

$$\begin{aligned} \int_{|E| \leq 3} |\tilde{n}(E) - n_{sc}(E)| dE &\leq CN^{-1/2+\varepsilon} + \int_{|E| \leq 3} |n(E) - n_{sc}(E)| dE \\ &\leq CN^{-1/2+\varepsilon} \end{aligned} \quad (4.22)$$

from (4.4). Thus we obtain from (4.21)

$$\frac{1}{N} \sum_{j=1}^N |\alpha_j - \gamma_j| \leq CN^{-1/2+\varepsilon}. \quad (4.23)$$

with an ε dependent constant.

To estimate $|\alpha_m - \gamma_m|$, we can assume, without loss of generality, that $\alpha_m \geq \gamma_m$, the other case is treated analogously. Let $\lambda > 0$ be a parameter that will be optimized later. Set $m_0 = \lceil CN\lambda^{3/2} \rceil$ with a sufficiently large constant C . Since $n_{sc}(-2 + \delta) \sim \delta^{3/2}$, the parameter λ is roughly the energy difference from the edge to the m_0 -th eigenvalue.

First we consider an index m such that $m_0 \leq m \leq N - m_0$. For a small positive number ℓ , define

$$S := \{j : \gamma_j \in [\gamma_m, \gamma_m + \ell]\}.$$

From the property $n_{sc}(-2 + \delta) \sim \delta^{3/2}$ for small δ , we have

$$|S| \geq cN\ell\lambda^{1/2}.$$

Now set $\ell = \min\{\frac{1}{2}|\alpha_m - \gamma_m|, c\lambda\}$ with some small positive constant c . Since for all $j \in S$

$$\alpha_j - \gamma_j \geq \alpha_m - (\gamma_m + \ell) \geq \frac{1}{2}(\alpha_m - \gamma_m) \geq \ell.$$

we have

$$\sum_{j=1}^N |\alpha_j - \gamma_j| \geq \sum_{j \in S} \ell \geq cN\ell^2\lambda^{1/2}.$$

Combining this estimate with (4.23), we have

$$\ell \leq C\lambda^{-1/4}N^{-1/4+\varepsilon/2}.$$

Assuming that $\lambda \geq CN^{-1/5+2\varepsilon/5}$, we see that $\ell = \frac{1}{2}|\alpha_m - \gamma_m|$ and we obtain

$$|\alpha_m - \gamma_m| \leq CN^{-1/5+\varepsilon} \quad (4.24)$$

for any m with $m_0 \leq m \leq N - m_0$.

For the extreme indices, we use that if $m \leq m_0$, then from Lemma 4.1 and (4.24), we have

$$-2 - CN^{-1/4+\delta} \leq \alpha_1 \leq \alpha_m \leq \alpha_{m_0} \leq \gamma_{m_0} + CN^{-1/5+\varepsilon} \leq -2 + C\lambda + CN^{-1/5+\varepsilon}$$

and

$$-2 \leq \gamma_m \leq -2 + C\lambda$$

for all $m \leq m_0$. Thus

$$|\alpha_m - \gamma_m| \leq C\lambda + CN^{-1/5+\varepsilon}$$

with C depending on ε . Similar estimates hold at the upper edge of the spectrum, i.e. for $m \geq N - m_0$. Choosing $\lambda = CN^{-1/5+\varepsilon}$, we conclude the proof of (4.18). The proof of (4.17) then follows from (4.7) and this concludes the proof of Proposition 4.4. $\square$

The following Proposition is a strengthening of the bound (4.23) used previously.

Proposition 4.5 *With the notations of Proposition 4.3, we have*

$$\frac{1}{N} \sum_i |\alpha_i - \gamma_i| \leq C\Phi + C\kappa N^{-1/2+\delta}. \quad (4.25)$$

Proof. We proceed similarly to the proof of Proposition 4.4 but we notice that in addition to (4.19), a stronger bound on $|n(E) - \tilde{n}(E)|$ is available for $E \in I := [E_-, E_+]$, where $E_{\pm} := \pm(2 - C_0\kappa)$, with some large constant C_0 . Here κ is subject to the constraints in (4.8). To obtain an improved bound, note that for any E in this interval

$$n(E) = \frac{1}{N} \sum_j \mathbb{E} \mathbf{1}(x_j \leq E) \leq \frac{1}{N} \sum_j \mathbb{E} \mathbf{1}(\alpha_j \leq E + \Phi) + Ce^{-cN^\delta} = \tilde{n}(E + \Phi) + Ce^{-cN^\delta}. \quad (4.26)$$

To see this inequality, define the random index

$$j_0 = j_0(E) := \max\{j : x_j \leq E\} = \sum_j \mathbf{1}(x_j \leq E)$$

and the deterministic index

$$j_1 = j_1(E) := \max\{j : \alpha_j \leq E + \Phi\} = \sum_j \mathbf{1}(\alpha_j \leq E + \Phi).$$

The estimate (4.26) will then follow if we prove that $j_0 \leq j_1$, i.e. $\alpha_{j_0} \leq E + \Phi$, with a very high probability. By (4.17) we have, with a very high probability, that

$$\gamma_{j_0} - CN^{-1/5+\delta} \leq x_{j_0} \leq E \leq x_{j_0+1} \leq \gamma_{j_0+1} + CN^{-1/5+\delta} \leq \gamma_{j_0} + CN^{-1/5+\delta}.$$

Therefore, with a very high probability, γ_{j_0} is in the $CN^{-1/5+\delta}$ vicinity of $E \in I$, and thus $CN\kappa^{3/2} \leq j_0 \leq CN(1 - \kappa^{3/2})$ holds for any fixed C if C_0 in the definition of $E_{\pm}$ is sufficiently large. Thus $|x_{j_0} - \alpha_{j_0}| \leq \Phi$ with a very high probability by (4.10), so $x_{j_0} \leq E$ implies $\alpha_{j_0} \leq E + \Phi$ and this proves (4.26).

The proof of the lower bound

$$n(E) \geq \tilde{n}(E - \Phi) - Ce^{-cN^\delta}$$

is analogous. Finally, by the Lipschitz continuity of $\tilde{n}(E)$ on a scale bigger than $(\log N)^2/N$, we have

$$|n(E) - \tilde{n}(E)| \leq C\Phi, \quad \forall E \in [E_-, E_+], \quad (4.27)$$

where we also used that $\Phi \geq C \exp(-cN^\delta)$.

Define the interval $J = [-2 - CN^{-1/4+\delta}, 2 + CN^{1/4+\delta}]$. Using (4.21), we have

$$\frac{1}{N} \sum_{j=1}^N |\alpha_j - \gamma_j| \leq (I) + (II) + (III) + \int |n(E) - n_{sc}(E)| dE \quad (4.28)$$

with

$$(I) := \int_I |\tilde{n}(E) - n(E)| dE, \quad (II) := \int_{J \setminus I} |\tilde{n}(E) - n(E)| dE, \quad (III) := \int_{J^c} |\tilde{n}(E) - n(E)| dE.$$

From (4.27) and (4.19), we have

$$(I) \leq C\Phi, \quad (II) \leq CN^{-1/2+\delta} |J \setminus I| \leq C\kappa N^{-1/2+\delta}$$

since $|J \setminus I| \leq C\kappa + N^{-1/4+\delta} \leq C\kappa$. Finally, $\tilde{n}(E) \equiv 0$ for $E < -2 - CN^{-1/4+\delta}$ and $\tilde{n}(E) \equiv 1$ for $E > 2 + CN^{-1/4+\delta}$ by (4.6). Combining these estimates with the fluctuation (4.7) and with the fact that $\mathbb{P}(\max |x_i| \geq K) \leq Ce^{-KN}$ for all large K , hence $n(E)(1 - n(E))$ decays exponentially for large $|E|$, we obtain

$$(III) \leq Ce^{-cN}.$$

Collecting all these estimates, inserting them into (4.28) and using (4.4), we conclude the proof of Proposition 4.5. $\square$

Proof of Lemma 2.2. We assume (2.21), (4.1), (4.2) and (4.4). We choose M, K, κ positive numbers satisfying (4.8) that will be optimized at the end of the proof.

Suppose now that $x_j \in I_j$ (recall the definition of I_j from (2.10)). Then, from the definition of b_j (2.15), we have

$$b_j = \frac{1}{N} \sum_{k: |k-j| > N\eta} \left[\frac{\text{sgn}(x_j - x_k)}{|x_j - x_k| + \eta} - \frac{\text{sgn}(x_j - \gamma_k)}{|x_j - \gamma_k| + \eta} \right] \quad (4.29)$$

Notice that function $g(x) := \frac{\text{sgn}(x)}{|x| + \eta}$ satisfies

$$|g(x) - g(y)| \leq \eta^{-2} |x - y| \quad (4.30)$$

as long as x and y have the same sign. In our case, $x_j - x_k$ and $x_j - \gamma_k$ have the same sign as long as $|x_k - \gamma_k| < |x_k - x_j|$. Applying the upper bound on the local density (4.1) for a grid of intervals of size $N^{-1+\delta}$, one can easily conclude that for any j, k with $|j - k| \geq N\eta$,

$$\mathbb{P}(|x_j - x_k| \leq \eta N^{-\delta}) \leq Ce^{-cN^{\delta/2}}$$

Combining this estimate with (4.17) and assuming

$$\eta \geq N^{-1/6},$$

we obtain that $x_j - x_k$ and $x_j - \gamma_k$ have the same sign apart from a set of very small probability. Applying (4.30), we have

$$\mathbb{P}\left(|b_j| \geq C\eta^{-2} \frac{1}{N} \sum_{|k-j| > N\eta} |x_k - \gamma_k|\right) \leq Ce^{-cN^{\delta/2}} \quad (4.31)$$

By (4.11) and (4.25), after recalling the definition of Φ (4.9), we have

$$\frac{1}{N} \sum_{|k-j|>N\eta} |x_k - \gamma_k| \leq \frac{1}{N} \sum_{|k-j|>N\eta} \left[|\alpha_k - \gamma_k| + |x_k - \alpha_k| \right] \leq C \left(\kappa N^{-1/2+\delta} + N^{-1/2+2\delta} K^{-1/2} + \frac{KM}{N} \right)$$

with a very high probability. Taking (4.31) into account, we can thus bound b_j by

$$|b_j| \leq C\eta^{-2} \left(\kappa N^{-1/2+\delta} + N^{-1/2+2\delta} K^{-1/2} + \frac{KM}{N} \right)$$

with a very high probability. On the complement event we estimate $|b_j|$ by $\eta^{-1} \leq N$. Recalling that $\mathbb{P} = \mathbb{P}_t$ denoted the probability with respect to $f_t \mu$, this gives

$$N \sum_j \int b_j^2 \mathbf{1}(x_j \in I_j) f_t d\mu_N \leq \eta^{-2} \Lambda$$

with

$$\Lambda := \eta^{-2} N^2 \left(\kappa N^{-1/2+\delta} + N^{-1/2+2\delta} K^{-1/2} + \frac{KM}{N} \right)^2.$$

Optimizing the parameters under the conditions (4.8) we can set

$$\kappa = N^{-1/18+\delta}, \quad M = N^{1/3}, \quad K = N^{1/9}$$

and finally $\Lambda \leq C\eta^{-2} N^{8/9+4\delta}$.

Suppose now that $x_j \notin I_j$. By (4.17) and $|b_j| \leq \eta^{-1} \leq N$ we have

$$N \sum_j \int b_j^2 \mathbf{1}(x_j \notin I_j) f_t d\mu \leq C e^{-cN^\delta} \quad (4.32)$$

This proves Lemma 2.2. □

A Some Properties of the Eigenvalue Process

In the main part of the paper we did not specify the function spaces in which the equations (2.6) and (3.1) are solved. In this appendix we summarize some basic properties of these equations. In particular, we justify the integration by parts in (3.6). For simplicity, we consider the most singular $\beta = 1$ case only.

The Dyson Brownian motion as a stochastic process was rigorously constructed in Section 4.3.1 of [14]. It was proved that the eigenvalues do not collide with probability one and thus (2.6) holds in a weak sense on the open set Σ_N . The coefficients of L have a $(x_i - x_j)^{-1}$ singularity near the coalescence hyperspace $x_i = x_j$. We focus only on the single collision singularities, i.e. on the case $j = i \pm 1$. By the ordering of the eigenvalues, higher order collision points form a zero measure set on the boundary of Σ_N and can thus be neglected. In an open neighborhood near the coalescence hyperspace $x_i = x_{i+1}$, the generator has the form

$$L = \frac{1}{2N} \left(\partial_i^2 + \partial_{i+1}^2 + \frac{1}{x_{i+1} - x_i} (\partial_{i+1} - \partial_i) \right) + L_{reg} = \frac{1}{4N} \left(\partial_v^2 + \partial_u^2 + \frac{1}{u} \partial_u \right) + L_{reg}, \quad u > 0,$$

after a change of variables, $v = \frac{1}{2}(x_i + x_{i+1})$, $u = \frac{1}{2}(x_{i+1} - x_i)$, where L_{reg} has regular coefficients. The boundary condition at $u = 0$ is given by the standard boundary condition of the generator of the Bessel process, $\partial_u^2 + \frac{1}{u}\partial_u$, which is $uf'(u) \rightarrow 0$ as $u \rightarrow 0+$. Thus L is defined on functions $f \in C^2(\Sigma_N)$ with sufficient decay at infinity and with boundary conditions

$$\lim_{x_{i+1}-x_i \rightarrow 0} (x_{i+1} - x_i)(\partial_{i+1} - \partial_i)f \rightarrow 0 \quad (\text{A.1})$$

for each i .

The generator $\tilde{L}$ of (3.1) differs from L only in drift terms with bounded coefficients, hence the boundary conditions of $\tilde{L}$ and L coincide. Finally, we need some non-vanishing and regularity property of the solution of (3.1):

Lemma A.1 *Let $\Omega \subset \Sigma^N$ be a bounded open set such that*

$$\overline{\Omega} \cap \bigcup_{i < j : (i,j) \neq (1,2)} \{\mathbf{x} : x_i = x_j\} = \emptyset$$

i.e. $\overline{\Omega}$ intersects at most one of the coalescent hyperplanes, namely the $\{x_1 = x_2\}$. Then any weak solution $q_t(\mathbf{x})$ of (3.1) with boundary conditions (A.1) is C^2 on $(t, \mathbf{x}) \in R_+ \times \overline{\Omega}$ and for any $t > 0$ we have

$$0 < \inf_{\overline{\Omega}} q_t \leq \sup_{\overline{\Omega}} q_t < \infty$$

Proof. The statement follows from regularity properties of the Bessel process with generator $\partial_u^2 + \frac{1}{u}\partial_u$. In a small neighborhood of the coalescence line $x_1 = x_2$ one can introduce a local coordinate system $(u, \mathbf{y}) = \Phi(\mathbf{x})$, where $u = \frac{1}{2}(x_2 - x_1) > 0$, $\mathbf{y} \in \mathbb{R}^{N-1}$, so that, in the case for GOE,

$$\tilde{L} = \frac{1}{4N} \left[\partial_u^2 + \frac{1}{u}\partial_u \right] + L_{reg},$$

where L_{reg} is an elliptic operator with second derivatives in the $\mathbf{y}$ variables and with bounded coefficients on the compact set $\Phi(\overline{\Omega})$. The solution in the new coordinates is $\tilde{q}_t(u, \mathbf{y}) = q_t(\Phi^{-1}(u, \mathbf{y}))$. Introducing a function $\hat{q}_t(a, b, \mathbf{y}) := \tilde{q}_t(\sqrt{a^2 + b^2}, \mathbf{y})$ defined in $N + 1$ variables, we see that $\hat{q}_t$ satisfies $\partial_t \hat{q}_t = \hat{L} \hat{q}_t$, where

$$\hat{L} = \frac{1}{N} \left[\partial_a^2 + \partial_b^2 \right] + L_{reg}.$$

i.e. $\hat{L}$ is elliptic with bounded coefficients in the new variables. Notice that the boundary condition (A.1) implies that, in the two dimensional plane of (a, b) , the support of the test function for the equation $\partial_t \hat{q}_t = \hat{L} \hat{q}_t$ is allowed to include the origin $(0, 0)$.

By standard parabolic regularity, we obtain that the solution is C^2 and is bounded from above and below. $\square$

This lemma justifies the integration by parts in (3.6). Since $q \in C^2$ and it is separated away from zero, $h = \sqrt{q}$ has no singularity on the coalescence lines. Since the function $\exp(-\tilde{\mathcal{H}})$ vanishes whenever $x_i = x_j$ for some $i \neq j$, the boundary terms of the form

$$\int_{x_i=x_j} \partial \sqrt{q} \partial^2 \sqrt{q} e^{-\tilde{\mathcal{H}}} d\mathbf{x}$$

in the integration by parts vanish.

B Logarithmic Sobolev inequality for convolution measures

Lemma B.1 Suppose K and H are two probability densities on $\mathbb{R}$ so that the LSI holds with constants a and b , respectively. Then LSI holds for their convolution $K * H$ as

$$\int_{\mathbb{R}} f(x) \log f(x) K * H(x) dx \leq \max(a, b) \int_{\mathbb{R}} (\nabla \sqrt{f(x)})^2 K * H(x) dx \quad (\text{B.1})$$

for any f with $\int f(x) K * H(x) dx = 1$. Here $\nabla = d/dx$.

Proof. The following proof is really a special case of the martingale approach used in [17] to prove LSI. Let

$$g(y) = \int_{\mathbb{R}} f(x) K(x - y) dx.$$

Then the left side of (B.1) is equal to

$$\int \int_{\mathbb{R}} \left[f(x) \log[f(x)/g(y)] + f(x) \log g(y) \right] K(x - y) dx H(y) dy. \quad (\text{B.2})$$

For any fixed y , from the LSI w.r.t. the measure $K(x - y) dx$, the first term on the right hand side is bounded by

$$a \int_{\mathbb{R}} \int_{\mathbb{R}} (\nabla \sqrt{f(x)})^2 K(x - y) dx H(y) dy.$$

Since $\int_{\mathbb{R}} gH = 1$, the second term in (B.2) is estimated by

$$\int_{\mathbb{R}} g(y) \log g(y) H(y) dy \leq b \int_{\mathbb{R}} (\nabla \sqrt{g(y)})^2 H(y) dy = \frac{b}{4} \int_{\mathbb{R}} g(y)^{-1} \left(\int_{\mathbb{R}} f(x) \nabla_y K(x - y) dx \right)^2 H(y) dy.$$

Integrating by parts, we can rewrite the last term as

$$\frac{b}{4} \int_{\mathbb{R}} g(y)^{-1} \left(\int_{\mathbb{R}} f'(x) K(x - y) dx \right)^2 H(y) dy \leq b \int_{\mathbb{R}} \int_{\mathbb{R}} (\nabla \sqrt{f(x)})^2 K(x - y) dx H(y) dy,$$

where we have used $f'(x) = 2\sqrt{f(x)}\nabla\sqrt{f(x)}$ and the Schwarz inequality. Combining these inequalities, we have proved the Lemma. $\square$

Acknowledgement: We thank Jun Yin for several helpful comments and pointing out some errors in the preliminary versions of this paper.

References

- [1] Bakry, D., Émery, M.: Diffusions hypercontractives. in: Séminaire de probabilités, XIX, 1983/84, **1123** Lecture Notes in Mathematics, Springer, Berlin, 1985, 177–206.
- [2] Ben Arous, G., Pécché, S.: Universality of local eigenvalue statistics for some sample covariance matrices. *Comm. Pure Appl. Math.* **LVIII**. (2005), 1–42.

- [3] Bleher, P., Its, A.: Semiclassical asymptotics of orthogonal polynomials, Riemann-Hilbert problem, and universality in the matrix model. *Ann. of Math.* **150** (1999): 185–266.
- [4] Brézin, E., Hikami, S.: Correlations of nearby levels induced by a random potential. *Nucl. Phys. B* **479** (1996), 697–706, and Spectral form factor in a random matrix theory. *Phys. Rev. E* **55** (1997), 4067–4083.
- [5] Deift, P., Kriecherbauer, T., McLaughlin, K.T-R, Venakides, S., Zhou, X.: Uniform asymptotics for polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory. *Comm. Pure Appl. Math.* **52** (1999):1335–1425.
- [6] Deift, P., Kriecherbauer, T., McLaughlin, K.T-R, Venakides, S., Zhou, X.: Strong asymptotics of orthogonal polynomials with respect to exponential weights. *Comm. Pure Appl. Math.* **52** (1999): 1491–1552.
- [7] Dyson, F.J.: A Brownian-motion model for the eigenvalues of a random matrix. *J. Math. Phys.* **3**, 1191–1198 (1962).
- [8] Erdős, L., Schlein, B., Yau, H.-T.: Semicircle law on short scales and delocalization of eigenvectors for Wigner random matrices. To appear in Ann. Probab. Preprint. arXiv.org:0711.1730
- [9] Erdős, L., Schlein, B., Yau, H.-T.: Wegner estimate and level repulsion for Wigner random matrices. To appear in Int. Math. Res. Notices (2008). Preprint arxiv.org/abs/0811.2591
- [10] Erdős, L., Ramirez, J., Schlein, B. and Yau, H.-T.: *Universality of sine-kernel for Wigner matrices with a small Gaussian perturbation*. Submitted to Electr. J. Prob. arxiv.org/abs/0905.2089
- [11] Erdős, L., Péché, S., Ramírez, J., Schlein, B. and Yau, H.-T.: Bulk universality for Wigner matrices. Submitted to Comm. Pure Appl. Math. Preprint arXiv.org:0905.4176.
- [12] Erdős, L., Ramírez, J., Schlein, B., Tao, T., Vu, V. and Yau, H.-T.: Bulk universality for Wigner hermitian matrices with subexponential decay. To appear in Math. Res. Letters. Preprint arXiv:0906.4400
- [13] Erdős, L., Yau, H.-T., Yin, J.: Universality for generalized Wigner matrices with Bernoulli distribution. Preprint arXiv:1003.3813
- [14] Guionnet, A.: Large random matrices: Lectures on Macroscopic Asymptotics. École d’Été de Probabilités de Saint-Flour XXXVI-2006. Springer.
- [15] Gustavsson, J.: Gaussian fluctuations of eigenvalues in the GUE, *Ann. Inst. H. Poincaré, Probab. Statist.* **41** (2005), no.2, 151–178
- [16] Johansson, K.: Universality of the local spacing distribution in certain ensembles of Hermitian Wigner matrices. *Comm. Math. Phys.* **215** (2001), no.3. 683–705.
- [17] S.L. Lu and H.T. Yau: Spectral gap and logarithmic Sobolev inequality for Kawasaki and Glauber dynamics, *Comm. Math. Phys.* **156**, 399–433, 1993.
- [18] Mehta, M.L.: Random Matrices. Academic Press, New York, 1991.

- [19] Pastur, L., Shcherbina M.: Bulk universality and related properties of Hermitian matrix models. *J. Stat. Phys.* **130** (2008), no.2., 205-250.
- [20] Sinai, Y. and Soshnikov, A.: A refinement of Wigner's semicircle law in a neighborhood of the spectrum edge. *Functional Anal. and Appl.* **32** (1998), no. 2, 114–131.
- [21] Soshnikov, A.: Universality at the edge of the spectrum in Wigner random matrices. *Comm. Math. Phys.* **207** (1999), no.3. 697-733.
- [22] Tao, T. and Vu, V.: Random matrices: Universality of the local eigenvalue statistics, *submitted*, Preprint arXiv:0906.0510.
- [23] Vu, V.: Spectral norm of random matrices. *Combinatorica*, **27** (6) (2007), 721-736.
- [24] Yau, H. T.: Relative entropy and the hydrodynamics of Ginzburg-Landau models, *Lett. Math. Phys.* **22** (1991) 63–80.